

CHE RUOLO HANNO GLI SCORE GERIATRICI ?

Malagola Michele, MD, PhD

Chair of Hematology, University of Brescia
Unit of Bone Marrow Transplant and Cellular Therapies
ASST Spedali Civili, Brescia



Corso di Formazione GITMO

Trapianto e terapie cellulari: cellule, donatori e condizionamenti


Udine, 8-9 Ottobre 2026

Aula Polifunzionale Ospedale di Udine


Disclosures of Name Surname

Company name	Research support	Employee	Consultant	Stockholder	Speakers bureau	Advisory board	Other
MEDAC					X		
Astellas					X		
MSD					X		



BACKGROUND (1)




Juliusson G. *Blood* 2012; 119: 3890-9; SEER. *Cancer Statistics Review* 1975-2008; Bhayat F et al. *BMC Cancer*. 2009;9:252..



BACKGROUND (2)





Transplantation and Cellular Therapy

journal homepage: www.tctjournal.org




Full Length Article
Allogeneic – Adult

GITMO Registry Study on Allogeneic Transplantation in Patients Aged ≥ 60 Years from 2000 to 2017: Improvements and Criticisms




Michele Malagola^{1,*}, Nicola Polverelli¹, Vicky Rubini¹, Massimo Martino², Francesca Patriarca³, Benedetto Bruno⁴, Luisa Giaccone⁴, Giovanni Grillo⁵, Stefania Bramanti⁶, Paolo Bernasconi⁷, Marco De Gobbi⁸, Annalisa Natale⁹, Elisabetta Terruzzi¹⁰, Attilio Olivieri¹¹, Patrizia Chiusolo¹², Angelo Michele Carella¹³, Marco Casini¹⁴, Chiara Nozzoli¹⁵, Patrizio Mazza¹⁶, Simona Bassi¹⁷, Francesco Onida¹⁸, Adriana Vacca¹⁹, Sadia Falcioni²⁰, Mario Luppi²¹, Anna Paola Iori²², Vincenzo Pavone²³, Cristina Skert²⁴, Paola Carluccio²⁵, Carlo Borghero²⁶, Anna Proia²⁷, Carmine Selleri²⁸, Nicoletta Sacchi²⁹, Sonia Mammoliti³⁰, Elena Oldani³¹, Fabio Ciceri³², Domenico Russo¹, Francesca Bonifazi³³



HCT-CI: is it useful in clinical practice above the age of 60 yrs?

Figure 4B.







Multidimensional geriatric assessment for elderly hematological patients (≥ 60 years) submitted to allogeneic stem cell transplantation. A French–Italian 10-year experience on 228 patients


2020

Nicola Polverelli ¹ · Paolo Tura^{1,2} · Giorgia Battipaglia ² · Michele Malagola¹ · Simona Bernardi ^{1,3} · Lisa Gandolfi¹ · Tatiana Zollner¹ · Camilla Zanaglio^{1,3} · Mirko Farina ¹ · Enrico Morello¹ · Alessandro Turra¹ · Mohamad Mohty² · Domenico Russo¹




Do you exclude a patient to allo-SCT or CART based on frailty?

Malagola M et al, Leuk & Lymph 2025



Do you exclude a patient to allo-SCT based on frailty?

Malagola M et al, Leuk & Lymph 2025



Udine, 8-9 Ottobre 2026


FRAILTY

- Frailty incidence in transplant setting: 10-25%
- Frailty is associated with poor outcome (morbidity and mortality)
- Dynamic process
- Can be reversible through pre-habilitation programs

Jayani R 2019; Klepin HD 2018; Williams GR 2014; Hamaker ME 2017; Muffly LS 2014; Deschler B 2018; Derman BA 2019; Holmes HM 2014; Lin RJ 2019; Nawas MT 2019; Salas MQ 2021

Multidimensional Geriatric Assessment

COMORBIDITY



FRAILITY and COMPREHENSIVE GERIATRIC ASSESSMENT

Ware JE 1992	- SF-36 PCS / SF-26 MCS	→ 36 item (QoL) for physical assessment 26 item (QoL) for mental assessment
Podsiadlo 1991	- TUG (Time Up and Go)	→ time for 3 m walking (two ways)
McDowell 1997	- ADL (Activities of Daily Life)	→ from 0 to 6 for daily living
Graf 2008	- IADL (Instrumental Activities of Daily Life)	→ from 0 to 8 for daily activities
Beelman 1997	- CES-D (Center for Epidemiologic Studies on Depression)	→ items for depression assessment
Saliba 2001	- VES13 (Vulnerable Elders Survey)	→ 13 items for geriatric assessment
Bland 2001	- 3-MS (Mini-Mental State Examination)	→ test for dementia
Fried 2001	- FFI (Fried Frailty Index)	→ 5 parameters (self reported and PS)
Rantanen 2003	- SPPB (Short Physical Performance Battery)	→ test for physical performance
Rockwood 2005	- CFS (Clinical Frailty Scale)	→ 9 point-scale (from very fit to terminal)
Rejesky 2008	- PAT-D (Pepper Assessment Tool for Disability)	→ 23 items on mobility+ADL+IADL
Hurria 2009	- DT (Distress Thermometer)	→ 1 item for level of distress (0 to 10)
Bellera 2012	- G8 (Geriatric 8)	→ 8 items (food, weight, mobility, BMI,...)
Bonanad 2015	- GAH scale (Geriatric Assessment in Hematology)	→ 30 items (medications, ADL, nutrition,..)
Palumbo 2015	- IMWG Frailty Score (Int Myeloma WG)	→ 30 item including CCI

Composite model	Risk groups		Outcomes at 2 years		Outcomes at 4 or 5 years	
	HCT-CI	KPS	NRM (%)	OS (%)	NRM (%)	OS (%)
Comorbidity/PS¹⁴ <i>Sorrer M Cancer 2008</i>	0-2	> 80%	16	68		
	0-2	≤ 80%	17	58		
	≥ 3	> 80%	30	41		
	≥ 3	≤ 80%	39	32		
Comorbidity/age score⁴⁶ (nonmyeloablative versus RIC) <i>Sorrer M JCO 2014</i>	HCT-CI/age					
	0		5-12	81-87		
	1-2		9-18	66-67		
	3-4		17-36	47-54		
	≥ 5		35-41	34-35		
Comorbidity/relapse score (patients ≥ 60 years old)⁴¹ <i>Sorrer M JAMA 2011</i>	HCT-CI	Relapse risk score				
	0	Low				69
	0	Standard				45
	0	High				41
<i>Elsawy MBBMT 2014</i>	1-2	Low				56
	1-2	Standard				44
	1-2	High				15
	≥ 3	Low				56
	≥ 3	Standard				23
	≥ 3	High				23
HCT-CI/EBMT⁴² <i>Muffy LS Haematol 2014</i>	HCT-CI	EBMT				
	0	< 4			11	72
	0	≥ 4			19	61
	1-2	< 4			16	63
	1-2	≥ 4			28	48
	≥ 3	< 4			31	40
	≥ 3	≥ 4			41	30
HCT-CI/IADL⁴³ HCT-CI score of ≥ 3 or IADL score < 14 acquire a score of 1. Both abnormalities get a score of 2	Scores					
	0			62		
	1			44		
	2			13		

Elsawy M et al, BMT 2016

Udine, 8-9 Ottobre 2026


Aula Polifunzionale Ospedale di Udine

Geriatric assessment to predict survival in older allogeneic hematopoietic cell transplantation recipients


Haematol 2014

Lori S. Muffy,¹ Masha Kocherginsky,² Wendy Stock,¹ Quynh Chu,¹ Michael R. Bishop,¹ Lucy A. Godley,¹ Justin Kline,¹ Hongtao Liu,¹ Olatoyosi M. Odenike,¹ Richard A. Larson,¹ Koen van Besien,³ and Andrew S. Artz¹


A



B



C



Udine, 8-9 Ottobre 2026

Corso di Formazione **GITMO** Trapianto e terapie cellulari: cellule, donatori e condizionamenti


Aula Polifunzionale Ospedale di Udine

Frailty assessment in adults undergoing allogeneic hematopoietic cell transplantation: insights from a multicenter GETH-TC study to optimize outcomes and care

Salas QM et al, Front Immun 2025


- 404 allo-SCT
- Age > 18 (median 56)
- 23% > 65 y

Study Design: Multicenter Implementation of the HCT Frailty Scale



Dynamics of Frailty According to Pre-Habilitation

Distribution of Patients Based on Pre-Habilitation	First Consultation N=62	HCT Admission N=62
PRE-HABILITATION N=61		
Fit	15 (24.2)	29 (46.8)
Pre-Frail	41 (66.1)	31 (50.0)
Frail	6 (9.7)	2 (3.2)
NO PRE-HABILITATION n=316	N=342	N=342
Fit	91 (26.6)	57 (16.7)
Pre-Frail	207 (60.5)	217 (63.5)
Frail	44 (12.9)	68 (19.9)
P VALUE	0.665	0.001






Udine, 8-9 Ottobre 2026




Corso di Formazione **GITMO** Trapianto e terapie cellulari: cellule, donatori e condizionamenti

Aula Polifunzionale Ospedale di Udine

HCT Frailty Scale at the First Consultation



HCT Frailty Scale at HCT Admission



HCT Pre-App program: a telemedicine-based prehabilitation for allogeneic hematopoietic cell transplant candidates

Blood Adv 2025

Raquel Sarahí Salinas Gonzalez,^{1,2} Laia Guardia,³ Berta Solé,³ Cristina Moreno,³ Noemi de Llobet,³ Enric Cascos,⁴ Julia Martinez-Sanchez,^{2,5,6} Maribel Díaz-Ricart,^{2,5,6} Enric Carreras,⁷ Maite Antonio,⁸ María Suárez-Lledó,^{2,3,6} Laura Rosiñol,^{2,3,6} Concepción Closa,¹ Raquel Sebio,¹ Carmen Martínez,^{2,3,6} Francesc Fernández-Avilés,^{2,3,6} Montserrat Rovira,^{2,3,6} and María Queralt Salas^{2,3,6}

NO PREHAB

n = 76

Frailty assessment only

PILOT PREHAB

n = 59

Individualized home exercise without digital supervision

TELE-PREHAB

n = 50

HCT Pre-App with remote rehabilitation supervision

HCT-FS

8 clinical, functional and laboratory variables



FIT • PREFRAIL • FRAIL

Assessment at consultation and HCT admission

HCT Pre-App program: a telemedicine-based prehabilitation for allogeneic hematopoietic cell transplant candidates

Blood Adv 2025


Raquel Sarahí Salinas Gonzalez,^{1,2} Laia Guardia,³ Berta Solé,³ Cristina Moreno,³ Noemi de Llobet,³ Enric Cascos,⁴ Julia Martinez-Sanchez,^{2,5,6} Maribel Díaz-Ricart,^{2,5,6} Enric Carreras,⁷ Maite Antonio,⁸ María Suárez-Lledó,^{2,3,6} Laura Rosiñol,^{2,3,6} Concepción Closa,¹ Raquel Sebio,¹ Carmen Martínez,^{2,3,6} Francesc Fernández-Avilés,^{2,3,6} Montserrat Rovira,^{2,3,6} and María Queralt Salas^{2,3,6}

HCT Pre-App prescription

Initial in-person functional assessment: 5-STs, 1-MSTs and DASI

3 structured sessions/week, at least 30 min each
Resistance and balance training, plus daily aerobic activity

Weekly digital follow-up and exercise adjustment until admission



Feasible and safe, with survival impact still uncertain

90.7%


recruited
59 of 65 eligible recipients

82.1%

overall exercise adherence

0

exercise-related adverse
events



CHARM is prognostic of geriatric morbidity and toxicity after allogeneic transplant for older adults: BMT CTN 1704 study

Blood Adv 2025

Andrew S. Artz,¹ Brent Logan,² Wael Saber,² Nancy Geller,³ Anna Bellach,³ Jianqun Kou,⁴ William A. Wood,⁵ John M. McCarty,⁶ Thomas G. Knight,⁷ Lyndsey Runaas,² Laura Johnston,⁸ Jeremy Walston,⁹ Ryotaro Nakamura,¹ Asmita Mishra,¹⁰ Joseph Uberti,¹¹ Parastoo B. Dahi,^{12,13} Jennifer N. Saultz,¹⁴ Shannon R. McCurdy,¹⁵ Lawrence Morris,¹⁶ Philip Imus,¹⁷ William J. Hogan,¹⁸ Kalyan Nadiminti,¹⁹ Vijaya Raj Bhatt,^{20,21} Rebecca Olin,²² Joseph Maakaron,²³ Ronald Sobecks,²⁴ Sarah Wall,²⁵ Deborah Mattila,⁴ Bailey Protz,⁴ Steven M. Devine,⁴ Mary M. Horowitz,² and Mohamed L. Sorrow^{26,27}

CHARM inputs **Prospective assessment**

1,105

Patients aged ≥60 years

49





US transplant centers



1 year

Prospective follow-up

Udine, 8-9 Ottobre 2026

Aula Polifunzionale Ospedale di Udine



A**B**

Artz et al, Blood Adv 2025

Article

Development of a Simplified Geriatric Score-4 (SGS-4) to Predict Outcomes After Allogeneic Hematopoietic Stem Cell Transplantation in Patients Aged over 50

2025

Eugenia Accorsi Buttini ^{1,2,*} , Alberto Zucchelli ^{1,3}, Paolo Tura ¹, Gianluca Bianco ¹ , Daniele Avenoso ^{1,4} , Giovanni Campisi ⁵, Mirko Farina ^{1,4} , Gabriele Magliano ⁴, Enrico Morello ⁴ , Vera Radici ^{1,4} , Nicola Polverelli ⁶ , Domenico Russo ^{1,4}, Alessandra Marengoni ^{1,3,†}  and Michele Malagola ^{1,4,†} 



Clinical Frailty Scale

Stato funzionale: ADL, IADL, Hand Grip, Gait Speed

Stato cognitivo: MMSE, MiniCOG


Screening: G8, VES-13, CFS, FIL score

Characteristics	N (%)
Number	135 (100)
Age at allo-SCT, years, median (range)	63 (50–75)
Male, n (%)	86 (63.7)
Diagnosis, n (%)	
AML	53 (39.3)
MDS	17 (12.6)
ALL	11 (8.1)
Lymphoma	13 (9.6)
MF	25 (18.5)
Other	16 (11.9)
DRI, n (%)	
Low	3 (2.2)
Intermediate	95 (70.4)
High	35 (25.9)
Very high	2 (1.5)


Characteristics	N (%)
KPS, n (%)	
90–100	126 (93.4)
70–80	9 (6.6)
HCT-CI, n (%)	
0	27 (20)
1–2	37 (27.4)
≥3	71 (52.6)
FIL score, n (%)	
Fit	132 (97.8)
Unfit	3 (2.2)
Disease status, n (%)	
1st CR	53 (39.3)
2nd or subsequent CR	82 (60.7)
Donor type, n (%)	
Related	33 (24.4)
Unrelated	63 (46.7)
Haploidentical	39 (28.9)
Stem cell source, n(%)	
PBSC	124 (91.9)
BM	11 (8.1)
Conditioning regimen, n (%)	
MAC	85 (63)
RIC	50 (37)
GVHD prophylaxis, n (%)	
CyA + MFF	3 (2.2)
CyA + MTX	4 (3)
CyA + MTX + ATG	78 (57.8)
PTCy + Other	27 (20)
Other	23 (17)
Follow-up, months, median (range)	7.5 (0.01–47.8)

Accorsi Buttini E, ..., Malagola M. Cancers 2025


OS




NRM



CIR




Accorsi Buttini E, ..., Malagola M. Cancers 2025




	Pts Evaluated	OS 1 y	OS 2 y	<i>p</i>	NRM 1y	NRM 2 y	<i>p</i>	CIR 1 y	CIR 2 y	<i>p</i>
ADL										
Any limitations	4	50	-		25	-		25	-	
No limitations	130	63	49	0.5	24	26	0.95	23	30	0.73
IADL										
Any limitations	118	30	50		46	24		35	35	
No limitations	15	66	30	0.11	21	46	0.11	22	21	0.72
G8, n										
≤13	51	52	46		24	24		31	37	
>13	83	67	48	0.7	24	29	0.9	17	26	0.23
VES 13, n										
≤3	14	11	11		30	30		58	58	
>3	120	67	52	0.04	23	26	0.93	18	28	0.01
FIL score										
FIT	132	65	50		24	27		21	30	
UNFIT	3	0	-	<0.01	0	23	0.41	100	-	<0.01
CFS										
FIT	75	63	55		23	23		22	35	
UNFIT	12	45	-	0.6	28	-	0.9	27	-	0.8
Gait Speed, m/s										
≤1.1	80	73	60		20	22		25	31	
>1.1	49	41	27	0.06	31	34	0.31	40	40	0.2
Chair Stand, times										
<14	90	65	50		21	22		25	32	
≥14	41	54	40	0.5	27	34	0.37	24	31	0.83
Hand Grip, Kg										
<26	37	45	30		34	38		28	41	
≥26	97	68	54	<0.01	21	21	0.05	20	28	0.6
Mini-Cog, n										
1-2	56	64	58		21	21		19	29	
3	77	60	39	0.28	24	29	0.44	26	31	0.45
MMSE, n										
≤24	13	19	19		52	52		29	48	
>24	121	66	50	0.03	22	24	0.2	22	29	0.56
HCT-CI, n										
0	27	88	33		17	17		18	25	
1-2	37	70	49		18	18		22	38	
≥3	71	55	37	0.2	29	34	0.2	28	30	0.88

Accorsi Buttini E, ..., Malagola M. Cancers 2025


OS ACCORDING TO HCT-CI



NRM ACCORDING TO Gait Speed




OS ACCORDING TO FIL




Accorsi Buttini E, ..., Malagola M. Cancers 2025

The Simplified Geriatric Score – SGS-4



Accorsi Buttini E, ..., Malagola M. *Cancers* 2025


The Simplified Geriatric Score – SGS-4



- Factor Analysis (Analisi dei Fattori Latenti)
- Calcolo dei Pesi delle Variabili sui Fattori Latenti (Metodo di Bartlett)
- Identificazione variabili con peso relativo > 0,4

Accorsi Buttini E, ..., Malagola M. *Cancers* 2025

The Simplified Geriatric Score – SGS-4




- Factor Analysis (Analisi dei Fattori Latenti)
- Calcolo dei Pesi delle Variabili sui Fattori Latenti (Metodo di Bartlett)
- Identificazione variabili con peso relativo > 0,4

Valori	Gait Speed	Hand Grip	Sesso
1	≤ 1.04	≤ 24.5	M
2	$1.04 < x \leq 1.17$	$24.5 < x \leq 32.7$	F
3	$1.17 < x \leq 1.29$	$32.7 < x \leq 41.3$	
4	$1.29 < x \leq 1.70$	$41.3 < x \leq 57.6$	

Accorsi Buttini E, ..., Malagola M. Cancers 2025

The Simplified Geriatric Score – SGS-4




- Factor Analysis (Analisi dei Fattori Latenti)
- Calcolo dei Pesi delle Variabili sui Fattori Latenti (Metodo di Bartlett)
- Identificazione variabili con peso relativo > 0,4

Valori	Gait Speed	Hand Grip	Sesso
1	≤ 1.04	≤ 24.5	M
2	$1.04 < x \leq 1.17$	$24.5 < x \leq 32.7$	F
3	$1.17 < x \leq 1.29$	$32.7 < x \leq 41.3$	
4	$1.29 < x \leq 1.70$	$41.3 < x \leq 57.6$	

Formula: [Gait Speed] + 2* [Hand Grip] + G8 + 1.5* [Sesso]

The Simplified Geriatric Score – SGS-4





- Factor Analysis (Analisi dei Fattori Latenti)
- Calcolo dei Pesi delle Variabili sui Fattori Latenti (Metodo di Bartlett)
- Identificazione variabili con peso relativo > 0,4

Valori	Gait Speed	Hand Grip	Sesso
1	≤ 1.04	≤ 24.5	M
2	$1.04 < x \leq 1.17$	$24.5 < x \leq 32.7$	F
3	$1.17 < x \leq 1.29$	$32.7 < x \leq 41.3$	
4	$1.29 < x \leq 1.70$	$41.3 < x \leq 57.6$	

Formula: [Gait Speed] + 2* [Hand Grip] + G8 + 1.5* [Sesso]

SCORE	≤ 13	FRAIL
	13 - 22.5	PRE-FRAIL
	> 22.5	FIT

Accorsi Buttini E, ..., Malagola M. Cancers 2025



Accorsi Buttini E, ..., Malagola M. Cancers 2025

Characteristics	Total N = 99	Frail N = 13	Prefrail N = 62	Fit N = 24	p-Value
Age, years (range)	63.0 (56.0, 66.0)	64.0 (56.0, 68.0)	62.0 (57.0, 66.0)	63.0 (56.0, 65.0)	>0.9
Diagnosis, n (%)					
AL	41 (41.4)	10 (76.9)	23 (37.1)	8 (33.3)	0.10
MDS	11 (11.1)	2 (15.4)	6 (9.7)	3 (12.5)	
Lymphoma	14 (14.1)	0 (0)	11 (17.7)	3 (12.5)	
MPN	1 (1.0)	0 (0)	0 (0)	1 (4.2)	
MM	5 (5.1)	0 (0)	5 (8.1)	0 (0)	
Other	27 (27.3)	1 (7.7)	17 (27.4)	9 (37.5)	
DRI, n (%)					
1	2 (2.0)	0 (0)	1 (1.6)	1 (4.2)	0.08
2	70 (70.7)	7 (53.8)	50 (80.7)	13 (54.2)	
3	24 (24.3)	6 (46.2)	10 (16.1)	8 (33.2)	
4	2 (2.0)	0 (0)	1 (1.6)	1 (4.2)	
missing	1 (1)	0	0	1 (4.2)	
KPS, n (%)					
90–100	92 (92.9)	11 (84.6)	57 (91.9)	24 (100)	0.6
70–80	7 (7.1)	2 (15.4)	5 (8.1)	0 (0)	
HCT-CI, n (%)					
0	21 (21.2)	1 (7.7)	12 (19.4)	8 (33.3)	0.13
1–2	28 (28.3)	3 (23.1)	16 (25.8)	9 (37.5)	
≥3	50 (50.5)	9 (69.2)	34 (54.8)	7 (29.2)	
Disease status, n (%)					
1st CR	33 (33.3)	6 (46.2)	20 (32.3)	7 (29.2)	0.6
2nd or subsequent CR	66 (66.7)	7 (53.8)	42 (67.7)	17 (70.8)	
Line of Therapy, n (%)					
1	48 (49.5)	8 (61.5)	28 (45.2)	12 (52.0)	0.5
≥2	50 (50.5)	5 (38.5)	34 (54.8)	11 (48.0)	
Missing	1 (1.0)	0	0	1	
Stem cell source, n (%)					
PBSC	91 (92.0)	12 (92.3)	56 (90.3)	23 (95.8)	0.9
BM	8 (8.0)	1 (7.7)	6 (9.7)	1 (4.2)	
Conditioning regimen, n (%)					
MAC	40 (40.4)	4 (30.7)	25 (40.3)	11 (45.8)	0.7
RIC	59 (59.6)	9 (69.3)	37 (59.7)	13 (54.2)	
TCI score, n	2.5	2.8	2.7	3.0	0.4
Donor age, years (range)	37 (28–55)	40 (28–45)	36 (28–47)	38 (28–55)	0.9
Donor, n (%)					
Related	23 (23.2)	3 (23.1)	13 (21.0)	7 (29.2)	0.9
Unrelated	44 (44.5)	6 (46.1)	27 (43.5)	11 (45.8)	
Haploidentical	32 (32.3)	4 (30.8)	22 (35.5)	6 (25.0)	
Time to allo-SCT, months (range)	11.3 (5.7–54.9)	6.3 (5.7–10.2)	12.4 (7.1–29.0)	19.5 (6.7–54.9)	0.03
aGVHD	48 (48.5)	4 (30.8)	29 (46.8)	15 (62.5)	0.17
cGVHD	16 (16.2)	0 (0.0)	10 (16.1)	6 (25.0)	0.12

Accorsi Buttini E, ..., Malagola M. Cancers 2025


Pre-Conditioning Frailty Phenotypes Influence Survival and Relapse for Older Allogeneic Transplantation Recipients

Sung AD et al, TCT 2025


Anthony D. Sung¹, Thuy Koll², Shannon H. Gier³, Alessandro Racioppi¹, Griffin White³, Meagan Lew¹, Marcia Free², Priyal Agarwal², Lauren M. Bohannon¹, Ernaya J. Johnson¹, Bharathi Selvan¹, Daria V. Babushok³, Noelle V. Frey³, Saar I. Gill³, Elizabeth O. Hexner³, MaryEllen Martin³, Alexander E. Perl³, Keith W. Pratz³, Selina M. Luger³, Nelson J. Chao¹, Alfred L. Fisher², Edward A. Stadtmauer³, David L. Porter³, Alison W. Loren³, Vijaya R. Bhatt², Phyllis A. Gimotty⁴, Shannon R. McCurdy³

- 280 allo-SCT
- Age > 60
- Fried's Frailty Phenotype (weight loss, exhaustion, slowness, weakness)
- Pre-admission or within 30 days
- 21 FRAIL, 161 PRE-FRAIL, 98 FIT

C. Patients aged 60-69 by Fried Frailty Phenotype




D. Patients Receiving RIC/NMA by Fried Frailty Phenotype




Udine, 8-9 Ottobre 2026

C. Non-relapse Mortality by Fried Frailty Phenotype



D. Relapse by Fried Frailty Phenotype




Sung AD et al, TCT 2025

Udine, 8-9 Ottobre 2026

Corso di Formazione **GITMO** Trapianto e terapie cellulari: cellule, donatori e condizionamenti

Aula Polifunzionale Ospedale di Udine

Aging in Mammals



Goertzel 2011

ALLO-GER Study

Multicentric Prospective Validation of the Simplified Geriatric Score (SGS-4) in Patients Aged ≥ 60 Years with Acute Leukemias (LA) or Myelodysplastic Syndrome (MDS) Undergoing Allogeneic Hematopoietic Stem Cell Transplantation (Allo-GER Study)

Principal Investigators:

Prof. Michele Malagola

Unit of Blood Diseases and Stem Cell Transplantation, Department of Clinical and Experimental Sciences, University of Brescia, ASST Spedali Civili, Brescia.

Co-Principal Investigators:

Prof. Alessandra Marengoni

Geriatric Unit, Department of Clinical and Experimental Sciences, University of Brescia, ASST Spedali Civili, Brescia.

Co-Investigators:

Dott. Luca Castagna

BMT Unit, AOR Villa Sofia Cervello, Palermo.

Prof. Alberto Zucchelli

Geriatric Unit, Department of Clinical and Experimental Sciences, University of Brescia, ASST Spedali Civili, Brescia.

Dott.ssa Eugenia Accorsi Buttini

PhD Artificial Intelligence in Medicine and Innovation In Clinical Research and Methodology, University of Brescia.

Dr Gianluca Bianco

PhD Artificial Intelligence in Medicine and Innovation In Clinical Research and Methodology, University of Brescia.

Prof Silvia Clara Giliani/ Dr.ssa Rosalba Monica Ferraro

Dep Molecular and Translational Medicine
University of Brescia

Endpoints	Primary endpoint • OS at 12 months after allo-SCT Secondary endpoints • NRM at 100 days, 6 months and 12 months after allo-SCT • CIR at 12 months after allo-SCT • Predictive value of cytokine patterns in combination with the SGS-4 score • Feasibility of the pre-transplant geriatric assessment, defined as the proportion of enrolled patients completing gait speed, handgrip strength, and G8 screening prior to allo-SCT.
Correlatives sub-studies	Two correlative biological sub-studies will be conducted in all the enrolled patients to elucidate the mechanisms underlying aging 3) Peripheral blood samples will be collected for the assessment of circulating biomarkers related to aging, frailty, and systemic inflammation. The biomarker panel will include markers of biological aging and inflammation (IL6, tumor necrosis factor receptor II, growth differentiation factor-15 [GDF-15], IGF1, insulin, cystatin-C). 4) Starting from 1.5ml of plasma, circulating extracellular vesicles (EVs) will be isolated using a commercial kit based on columns that isolate small EVs by affinity. Then, Fourier transform infrared spectroscopy (FT-IR) will be performed using a Cary 630 FT-IR Spectrometer (Agilent Technologies) operating in transmission mode. From each sample, a volume of 2 µL of EV suspension or 2 µL of whole plasma, depending on the analysis. FT-IR spectroscopy will simultaneously provide information on the main classes of biomolecules: lipids, proteins, nucleic acids, and carbohydrates, allowing the chemical-structural composition and organization of cellular systems and EVs to be investigated.

Statistical design

Sample size calculation.

A total of approximately 300 patients will be enrolled in this study. This sample size was estimated assuming a 1-year mortality rate of 30%, corresponding to ~100 events—sufficient for robust model validation. With this size, the 95% CI around an expected AUC of 0.65 is estimated at ± 0.07 . For FT-IR spectra analyses, subsequent bioinformatic and biostatistics analyses (PCA and LDA) will be necessary to compare the different components and identify biological subgroups among the samples analyzed.

2 years for recruitment

1 year of follow up for the last recruited patient

1. TMO Adulti - ASST Spedali Civili, Brescia.
Michele Malagola - michele.malagola@unibs.it
2. SC Ematologia – IRCCS Fondazione Policlinico San Matteo, Pavia.
Nicola Polverelli - n.polverelli@smatteo.pv
3. UOC Trapianto e Terapie Cellulari in Ematologia – IRCCS AOU di Bologna, Bologna.
Francesca Bonifazi – crc-ematologia@aosp.bo.it
4. UOC Ematologia e Trapianto di Cellule staminali emopoietiche – Fondazione Policlinico Universitario Agostino Gemelli IRCCS
Patrizia Chiusolo - patrizia.chiusolo@unicatt.it
5. Unità Trapianti di Midollo Osseo e Terapie Cellulari – AOR Villa Sofia Cervello, Palermo
Luca Catsagna - carmelocostanza85@gmail.com – l.castagna@villasofia.it
6. UOSC Ematologia con Trapianto CSE - AORN A. Cardarelli , Napoli
Alessandra Picardi - alessandra.picardi@aocardarelli.it
7. Clinica di Ematologia,Trapianto e Terapia Cellulare Azienda Ospedaliero - Universitaria delle Marche
Francesco Saraceni – irene.federici@ospedaliriuniti.marche.it
8. SS Centro Trapianti di Midollo e Terapie Cellulari - Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico di Milano
Giorgia Saporiti – giulia.milesi@policlinico.mi.it
9. Divisione di Ematologia - Azienda Ospedaliera S. Croce e Carle, Cuneo
Nicola Mordini - mordini.n@ospedale.cuneo.it
10. SODc Ematologia - AOU Careggi, Firenze
Chiara Nozzoli - ctc.smo@aou-careggi.toscana.it
11. UOC Ematologia e Unità Terapia Intensiva Ematologica – Casa sollievo della Sofferenza IRCCS, San Giovanni Rotondo
Angelo Michele Carella - am.carella@operapadrepio.it
12. UO Ematologia – IRCCS Ospedale San Raffaele, Milano
Maria Teresa Lupo Stanghellini – regolatorio.scp@hsr.it
13. Struttura Complessa a Direzione Universitaria di Ematologia - Azienda Ospedaliera Ss Antonio, Biagio e Cesare Arrigo, Alessandria
Lucia Brunello – clinicaltrialcenter@ospedale.al.it
14. Clinica Ematologica e Unità di Terapie Cellulari - Azienda Sanitaria Universitaria Friuli Centrale, Udine
Francesca Patriarca - francesca.patriarca@asufc.sanita.fvg.it
15. UO CTMO – Grande Ospedale Metropolitano “Bianchi-Melacrino-Morelli”, Reggio Calabria
Massimo Martino – ctc@ospedalerc.it
16. UOC Ematologia e Trapianto di Midollo - AOU Federico II, Napoli
Giorgia Battipaglia – giorgia.battipaglia@unin.it
17. SS Trapianto Allogeneico di Cellule Staminali, Azienda Ospedaliero-Universitaria Città della Salute e della Scienza di Torino, Torino
Alessandro Busca – abusca@cittadellasalute.to.it

CET Submission – 30/04/26 / 12/6/26

CET Approval - 5/8/26

ASST Spedali Civili Authorisation – 15/9/26

1° Patient (BS) – 21/9/26

Centers' activation ongoing

Frailty assessment and allo-SCT

- Allo-SCT may cure approximately 50% of elderly patients (in particularly AL > 60 yrs)
- Age is not a “stand alone” selection criteria
- Comorbidities are not enough → modulation of transplant platforms
- **PATIENT'S SELECTION IS CRUCIAL**
- Different scores? Is there a winner?
- Underlying biological mechanisms
- Assessment of vulnerabilities (pre-habilitation) → **PROSPECTIVE STUDIES !!**

